

KAN REPLACEMENT OF SIMPLICIAL MANIFOLDS


CHENCHANG ZHU

COURANT RESEARCH CENTRE “HIGHER ORDER STRUCTURES”, UNIVERSITY OF
GÖTTINGEN

ABSTRACT. We establish a functor Kan from local Kan simplicial manifolds to weak Kan simplicial manifolds. It gives a solution to the problem of extending local Lie groupoids to Lie 2-groupoids.

1. INTRODUCTION

It is a classical topic to study the correspondence between global and infinitesimal symmetries. For us, the process from global symmetries to infinitesimal ones is called differentiation, and the inverse process is called integration. A classical example of such is in the case of Lie groups and Lie algebras,



However, when our symmetries become more complicated, such as L_∞ -algebras, or even L_∞ -algebroids, the integration and differentiation both become harder. The following problems have been solved for these higher symmetries: integration of nilpotent L_∞ -algebras by Getzler [5], integration of general L_∞ -algebras by Henriques [6], differentiation of L_∞ -groupoids by Ševera [13], both directions for Lie 1-algebroids by Cattaneo-Felder [2], Crainic-Fernandes [3], and from a higher viewpoint by Tseng-Zhu [11]. Here the author wants to emphasis a middle step of local symmetries missing in the above correspondence,



Indeed, to do differentiation to obtain infinitesimal symmetries, we only need local symmetries. Conversely, sometimes, it is easier to obtain a local integration, avoiding some analytic issues (for example in [5] for L_∞ -algebras). In this paper, we make our first attempts towards the extension problem from local symmetries to global ones: we construct an extension from local Kan simplicial manifolds to weak Kan ones. The classical extension of local Lie group to a topological group discussed by van Est in [12] can be viewed as 1-truncation of our result. Its 2-truncation applied to local Lie groupoids provide a solution to the integration problem of Lie algebroids to Lie 2-groupoids [15]. Notice that unlike Lie algebras which one-to-one correspond to simply connected Lie groups, Lie algebroids (integrable or not) one-to-one correspond to a sort of Lie 2-groupoids with some étale property.

We use the viewpoint of Kan simplicial manifolds to describe arbitrary Lie n -groupoids.

Date: September 3, 2009.

Supported by the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG)) through the Institutional Strategy of the University of Göttingen.


Recall that a simplicial manifold X consists of manifolds X_n and structure maps

$$(1) \quad d_i^n : X_n \rightarrow X_{n-1} \text{ (face maps)} \quad s_i^n : X_n \rightarrow X_{n+1} \text{ (degeneracy maps), for } i \in \{0, 1, 2, \dots, n\}$$

that satisfy suitable coherence conditions (see for example [4]). The first two examples of simplicial manifolds (actually, they are simplicial sets with discrete topology) are the simplicial m -simplex $\Delta[m]$ and the horn $\Lambda[m, j]$ with

$$(2) \quad \begin{aligned} (\Delta[m])_n &= \{f : (0, 1, \dots, n) \rightarrow (0, 1, \dots, m) \mid f(i) \leq f(j), \forall i \leq j\}, \\ (\Lambda[m, j])_n &= \{f \in (\Delta[m])_n \mid \{0, \dots, j-1, j+1, \dots, m\} \not\subseteq \{f(0), \dots, f(n)\}\}. \end{aligned}$$

The horn $\Lambda[m, j]$ should be thought as a simplicial set obtained from $\Delta[m]$ by taking away its unique non-degenerate m -simplex as well as the j -th of its $m+1$ non-degenerate $(m-1)$ -simplices.



Our convention for arrows is that they are oriented from bigger numbers to smaller numbers.

Let us recall that in homotopy theory, Kan conditions say that the natural restriction map

$$(3) \quad X_m = \text{hom}(\Delta[m], X) \rightarrow \text{hom}(\Lambda[m, j], X).$$

is surjective, i.e. any horn can be filled up by a simplex. They correspond to the possibility of composing and inverting various morphisms, in the language of groupoids.

With enrichment in differential geometry, **Kan conditions** are

$$Kan(m, j): (3) \text{ is a surjective submersion,} \quad Kan^!(m, j): (3) \text{ is a diffeomorphism.}$$

But since $\text{hom}(\Lambda[m, j], X)$ is formed by taking a numerous fibre products of the X_i 's, it may not be a manifold. However if (3) is a submersion for all $0 \leq j \leq m \leq m_0$, as shown in [6, Lemma 2.4], $\text{hom}(\Lambda[m_0, j], X)$ is a manifold for all $0 \leq j \leq m_0$. Hence we are allowed to define,

Definition 1.1. A *Lie n -groupoid* X ($n \in \mathbb{N} \cup \infty$) is a simplicial manifold that satisfies $Kan(m, j)$, $\forall m \geq 1$, $0 \leq j \leq m$, and $Kan^!(m, j)$, $\forall m > n$, $0 \leq j \leq m$. When $n = \infty$, a Lie ∞ -groupoid is also called a *Kan simplicial manifold*.

Then Lie 1-groupoid is simply the nerve of a Lie groupoid.

To describe local Lie groupoids, we need **local Kan conditions**:

$$Kan^l(m, j): (3) \text{ is a submersion,} \quad Kan^{!l}(m, j): (3) \text{ is injective étale.}$$

Definition 1.2. A *local Lie n -groupoid* X ($n \in \mathbb{N} \cup \infty$) is a simplicial manifold that satisfies $Kan^l(m, j)$, $\forall m \geq 1$, $0 \leq j \leq m$, and $Kan^{!l}(m, j)$, $\forall m > n$, $0 \leq j \leq m$. When $n = \infty$, a local Lie ∞ -groupoid is also called a *local Kan simplicial manifold*.

Then a local Lie 1-groupoid X is the nerve of a local Lie groupoid.

As soon as we have done this, it becomes clear that to associate a Kan object $Kan(X)$ to a local Kan simplicial manifold X , we need to do some sort of fibrant replacement in

A calculation shows that

$$\begin{aligned}
X_2^1 &= X_2 \sqcup X_1 \times_{X_0} X_1 \sqcup X_1 \times_{X_0} X_1 \sqcup X_1 \times_{X_0} X_1 \\
&\quad \sqcup (\sqcup_{j=0}^3 \text{hom}(\Lambda[3, j], X)) \\
X_2^2 &= X_2^1 \sqcup X_1^1 \times_{X_0} X_1^1 \sqcup X_1^1 \times_{X_0} X_1^1 \sqcup X_1^1 \times_{X_0} X_1^1 \\
&\quad \sqcup (\sqcup_{j=0}^3 \text{hom}(\Lambda[3, j], X^1)) \\
&\quad \vdots
\end{aligned}$$

Inside X_2^1 , there are three copies of $X_1 \times_{X_0} X_1$. The first is an artificial filling of the horn $X_1 \times_{X_0} X_1$, and the second two are images of degeneracies of $X_1 \times_{X_0} X_1$ in X_1^1 . The same for X_2^2 , etc. We represent an element in X_2^1 as

$$(8) \quad X_2 : \triangle, \quad X_1 \times_{X_0} X_1 : \text{---}, \quad \text{hom}(\Lambda(3, j), X) : \triangle \dots 4 \text{ such}$$

plus those degenerate ones in the other two copies of $X_1 \times_{X_0} X_1$. Furthermore we represent an element in X_2^2 as

X_2^1 : described as above

$$X_1^1 \times_{X_0} X_1^1 : \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

$$(9) \quad \text{hom}(\Lambda[3, j], X^1) : \begin{array}{c} \triangle \\ \triangle \\ \triangle \end{array} \quad \triangle \quad \triangle \quad \triangle \quad \triangle \quad \dots$$

We now show that

$$(10) \quad X_2^1 \rightarrow \text{hom}(\Lambda[2, 0], X^1) = X_1^1 \times_{d_2, X_0, d_1} X_1^1$$

is actually not a submersion.

We first need some technical preparation. A simplicial set S is *collapsible* if it admits a filtration

$$(11) \quad pt = S_0 \subset S_1 \subset \dots \subset S_k = S$$

such that each S_i is obtained from the previous one by filling a horn, namely such that S_i can be written as $S_i = S_{i-1} \sqcup_{\Lambda[n_i, l_i]} \Delta[n_i]$ for some injective map $\Lambda[n_i, l_i] \hookrightarrow S_{i-1}$. Thus we have an order for collapsible simplicial sets: we say S is not bigger than T , denoted as $S \prec T$, if $T = S_t$ and $S = S_s$ with $s \leq t$ in (11). For us the notation $S \prec T$ also indicates the inclusion map $S \rightarrow T$. We also define the dimension of a collapsible simplicial set S as

$$\dim S = \max_k \{\Delta[k] \prec S\}.$$

Lemma 2.1. *Given $S \prec T$ and a local Kan simplicial manifold X , $\text{hom}(T, X)$ and $\text{hom}(S, X)$ are both manifolds, and the natural map*

$$\text{hom}(T, X) \rightarrow \text{hom}(S, X),$$

is always a submersion.

This is proven in [6, Lemma 2.4] for Kan simplicial manifolds, but it is easy to see that it works also for local Kan ones since only the submersion condition is used.

Back to the map (10), X_2^1 has several components, and the horn projection map (10) induces on each component the following maps:

$$\begin{aligned} (12) \quad & X_2 \rightarrow \text{hom}(\Lambda[2, 0], X), & \text{hom}(\Lambda[3, j], X) &\rightarrow \text{hom}(\Lambda[2, 0], X) \\ (13) \quad & X_1 \times_{X_0} X_1 \rightarrow X_1 \times_{X_0} (X_1 \times_{X_0} X_1), & X_1 \times_{X_0} X_1 &\rightarrow (X_1 \times_{X_0} X_1) \times_{X_0} (X_1 \times_{X_0} X_1), \\ (14) \quad & X_1 \times_{X_0} X_1 \rightarrow X_1 \times_{X_0} X_1. \end{aligned}$$

The morphisms in (12) are always submersions by Lemma 2.1. However the morphisms in (13) are not submersions. This implies that $\text{hom}(\Delta[2], \text{Kan}(X)) \rightarrow \text{hom}(\Lambda[2, 0], \text{Kan}(X))$ will not be a submersion. Hence $\text{Kan}(X)$ will not be a Kan simplicial manifold, but we will prove that it satisfies

- (A) $\text{Kan}(X)$ is a simplicial manifold;
- (B) moreover, $\text{hom}(S, \text{Kan}(X))$ is a manifold for any collapsible S ;
- (C) the natural map $\text{Kan}(X)_n \rightarrow \text{hom}(\Lambda[n, l], \text{Kan}(X))$ is surjective for all n and l with $0 \leq l \leq n$.

Simplicial manifolds satisfying such conditions are called **weak Kan simplicial manifolds**. These weak Kan simplicial manifolds as simplicial sets are indeed Kan. The submersion condition in Kan condition is replaced by condition (B). By Lemma 2.1, we can see that the submersion condition implies (B), hence weak Kan is indeed weaker than Kan. However, the usual place to use submersion condition is to guarantee some representibility for example the one in condition (B). Hence we see that in many cases we can bypass the difficulty thanks to this condition. For example, we can still talk about hypercovers of these weak Kan simplicial manifolds (even though it is not used in this paper).

Before attacking the problem, we first prepare a technical lemma:

Lemma 2.2. *Suppose that X^β 's are a sequence of simplicial sets constructed by (6),*

- (1) *if S is a collapsible simplicial set, then we can decompose*

$$\text{hom}(S, X^{\beta+1}) = \sqcup_{a \in A} \text{hom}(S_a, X^\beta),$$

with a finite set of collapsible simplicial sets $\{S_a : a \in A\} \ni S$ satisfying $\dim S_a \leq \dim S$.

- (2) *this decomposition respects morphisms, that is if $S \prec T$, and both of them have a decomposition,*

$$\text{hom}(S, X^{\beta+1}) = \sqcup_{a \in A} \text{hom}(S_a, X^\beta), \quad \text{hom}(T, X^{\beta+1}) = \sqcup_{a' \in A'} \text{hom}(T_{a'}, X^\beta),$$

then there is a map $a : A' \rightarrow A$, and morphisms of simplicial sets $S_{a(a')} \rightarrow T_{a'}$, such that the natural morphism $\text{hom}(T, X^{\beta+1}) \rightarrow \text{hom}(S, X^{\beta+1})$ is induced from $\text{hom}(T_{a'}, X^\beta) \rightarrow \text{hom}(S_{a(a')}, X^\beta)$ on the level of their decompositions.

Proof. Since the procedure to form $X^{\beta+1}$ by X^β is the same as the one to form X^1 by X , we only have to prove the two statements for $\beta = 0$. Since X is arbitrary, the decomposition in (1) is clearly unique. We use an induction on the size of S and T . The initial assumption is verified in the calculation we did earlier in this section. It is clear that (1) holds for $\Delta[m]$ for $m \in \mathbb{N}$. Now with a fixed n , we consider a horn filling diagram as we mentioned in the

process of (11)

$$(15) \quad \begin{array}{ccc} S & \xrightarrow{\quad} & T \\ \uparrow & & \downarrow \\ \Lambda[k, j] & \xrightarrow{\quad} & \Delta[k] \end{array} ,$$

with $\dim S \leq \dim T \leq n-1$, and $k \leq n-1$.

We suppose that

- (i) statement (1) is true for all S' with $S' \prec S$;
- (ii) statement (2) is true for $S' \prec T'$ and $S' \prec \Delta[n]$ when $\dim S' \leq \dim T' \leq n-1$ and when (1) verifies for S' and T' .

To finish the induction, we will prove that

- statement (1) holds for T ;
- statement (2) holds for $S \prec T$, and $T \prec \Delta[n]$ if such a map $T \rightarrow \Delta[n]$ exists.

First of all, we apply $\text{hom}(-, X^1)$ to (15) and apply the induction hypothesis to $\text{hom}(S, X^1)$, $\text{hom}(\Lambda[k, j], X^1)$ and $\text{hom}(\Delta[k], X^1)$, then we have

$$\begin{aligned} \text{hom}(T, X^1) &= \text{hom}(S, X^1) \times_{\text{hom}(\Lambda[k, j], X^1)} \text{hom}(\Delta[k], X^1) \\ &= \sqcup_{a' \in A'} \text{hom}(S'_{a'}, X) \times_{\sqcup_{a \in A} \text{hom}(S_a, X)} \sqcup_{a'' \in A''} \text{hom}(S''_{a''}, X) \\ &= \sqcup_{b \in B} \text{hom}(T_b, X). \end{aligned}$$

Here T_b is formed when $a(a') = a(a'')$ by

$$\begin{array}{ccc} S'_{a'} & \xrightarrow{\quad} & T_b \\ \uparrow & & \downarrow \\ S_{a(a')} & \xrightarrow{\quad} & S''_{a''} \end{array} .$$

We obtain a map $B \rightarrow A'$ defined by $b \mapsto a'$ and morphisms $S'_{a'} \rightarrow T_b$. They induce the morphisms $\text{hom}(T_b, X) \rightarrow \text{hom}(S'_{a'}, X)$, hence the morphism $\text{hom}(T, X^1) \rightarrow \text{hom}(S, X^1)$. It's not hard to see that $T \in \{T_b\}$ by induction hypothesis and (15).

Suppose $\text{hom}(\Delta[n], X^1) = \sqcup_{c \in C} (D_c, X)$. If there is a map $T \prec \Delta[n]$, by restriction, we obtain maps $S \prec \Delta[n]$, $\Lambda[k, j] \prec \Delta[n]$, and $\Delta[k] \prec \Delta[n]$ which fit in the following commutative diagram:

$$\begin{array}{ccc} S & \xrightarrow{\quad} & \Delta[n] \\ \uparrow & \nearrow & \uparrow \\ \Lambda[k, j] & \xrightarrow{\quad} & \Delta[k] \end{array} ,$$

By induction hypothesis, we have

- the morphism $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(S, X^1)$ is induced by a map $a' : C \rightarrow A'$ and morphisms $S'_{a'(c)} \rightarrow D_c$;
- the morphism $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(\Lambda[k, j], X^1)$ is induced by a map $a : C \rightarrow A$ and morphisms $S_{a(c)} \rightarrow D_c$;

- the morphism $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(\Delta[k], X^1)$ is induced by a map $a'' : C \rightarrow A''$ and morphisms $S''_{a''(c)} \rightarrow D_c$.

We see that $\text{hom}(D_c, X) \rightarrow \text{hom}(S_{a(c)}, X)$ induces $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(\Delta[k, j], X^1)$, and the composed morphism $\text{hom}(D_c, X) \rightarrow \text{hom}(S'_{a'(c)}, X) \rightarrow \text{hom}(S_{a(a'(c))}, X)$ induces $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(S, X^1) \rightarrow \text{hom}(\Delta[k, j], X^1)$, which is the same morphism as $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(\Delta[k, j], X^1)$. Hence by uniqueness of the decomposition, we have $a(a'(c)) = a(c)$ and similarly $a(c) = a(a''(c))$, and a commutative diagram

$$\begin{array}{ccc} S'_{a'(c)} & \xrightarrow{\quad} & D_c \\ \uparrow & \nearrow & \uparrow \\ S_{a(a')} & \xrightarrow{\quad} & S''_{a''(c)} \end{array} \quad ,$$

Then $T_{b(c)}$ defined by the pushout diagram

$$\begin{array}{ccc} S''_{a''(c)} & \xrightarrow{\quad} & T_{b(c)} \\ \uparrow & \lrcorner & \uparrow \\ S_{a(c)} & \xrightarrow{\quad} & S'_{a'(c)} \end{array} \quad .$$

has a canonical map $T_{b(c)} \rightarrow D_c$. By the property of $\text{hom}(T, X^1)$ being the fibre product, these canonical maps induce the map $\text{hom}(\Delta[n], X^1) \rightarrow \text{hom}(T, X^1)$ via the maps $\text{hom}(D_c, X) \rightarrow \text{hom}(T_{b(c)}, X)$. $\square$

Proposition-Definition 2.3. The operation Kan constructed in (6) is a functor from the category of local Kan manifolds X to the one of weak Kan simplicial manifolds.

Proof. The construction of Kan makes it clear that it is functorial. Since $Kan(X) = \text{colim}_\beta X^\beta$, given any finite simplicial set A (a collapsible simplicial set S is such), the natural map of sets is an isomorphism,

$$(16) \quad \text{colim}_\beta \text{hom}(A, X^\beta) \xrightarrow{\cong} \text{hom}(A, Kan(X)).$$

Moreover by Lemma 2.2,

$$(17) \quad \text{hom}(S, X^{\beta+1}) = \text{hom}(S, X^\beta) \bigsqcup (\bigsqcup_a \text{hom}(S_a, X^\beta)),$$

We then use Lemma 2.2 recursively, and obtain that for any collapsible simplicial set T ,

$$\text{hom}(T, X^\beta) = \bigsqcup \text{hom}(T_p, X),$$

for a finite set of collapsible simplicial sets T_p . Hence $\text{hom}(S, X^\beta)$ and $\text{hom}(S_a, X^\beta)$ are manifolds because X is local Kan. By (17) and (16), $\text{hom}(S, Kan(X))$ is a disjoint union of manifolds.

So it remains to show that $Kan(X)$ is Kan as a simplicial set. We take an element $A \rightarrow B$ of J and a solid arrow diagram,

$$(18) \quad \begin{array}{ccc} A & \longrightarrow & Kan(X) \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ B & \longrightarrow & pt \end{array}$$

then we must show that the dotted arrow exists. By the isomorphism (16), the map $A \rightarrow Kan(X)$ factors through $X^\beta \rightarrow Kan(X)$ for some β and we have the solid arrow diagram

$$\begin{array}{ccccccc} A & \longrightarrow & X^\beta & \longrightarrow & X^{\beta+1} & \longrightarrow & Kan(X) \\ \downarrow & & \downarrow & \nearrow \text{dotted} & \searrow & \nearrow & \\ B & \longrightarrow & pt & & & & \end{array}$$

Since $X^{\beta+1}$ is constructed as the push-out in (6), the dotted arrow naturally exists, and this dotted arrow defines the one in (18).

Now we only have to verify that the dotted arrow in (18) exists for $\Lambda[1, j] \rightarrow \Delta[1]$ for $j = 0, 1$ and $\Lambda[2, j] \rightarrow \Delta[2]$ for $j = 0, 2$. We have $X_0^\beta = X_0$, and

$$\text{hom}(\Lambda[1, j], X^{\beta+1}) = X_0^{\beta+1} = X_0, \quad \text{hom}(\Delta[1], X^{\beta+1}) = X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta,$$

thus the map $\text{hom}(\Delta[1], X^{\beta+1}) \rightarrow \text{hom}(\Lambda[1, j], X^{\beta+1})$ being the pull-back of d_1 or d_0 , has to be a surjective submersion. Now we prove that if X^β is invertible, then $X^{\beta+1}$ is also invertible.

$$\begin{aligned} & \text{hom}(\Lambda[2, 2], X^{\beta+1}) \\ &= X_1^{\beta+1} \times_{d_1, X_0, d_1} X_1^{\beta+1} \\ &= (X_1^\beta \sqcup X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \times_{d_1, X_0, d_1} (X_1^\beta \sqcup X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \\ &= X_1^\beta \times_{d_1, X_0, d_1} X_1^\beta \sqcup X_1^\beta \times_{d_1, X_0, d_1} (X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \sqcup (X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \times_{d_1, X_0, d_1} X_1^\beta \sqcup \dots \end{aligned}$$

Since X^β is invertible, $X_1^\beta \times_{d_1, X_0, d_1} X_1^\beta \cong X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta$. Hence

$$\begin{aligned} & X_1^\beta \times_{d_1, X_0, d_1} (X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \cong X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta, \\ & (X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \times_{d_1, X_0, d_1} X_1^\beta \cong X_1^\beta \times_{d_1, X_0, d_1} X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta \cong (X_1^\beta \times_{d_0, X_0, d_1} X_1^\beta) \times_{d_0, X_0, d_1} X_1^\beta, \\ & \dots \end{aligned}$$

It is easy to continue to verify that $X^{\beta+1}$ is invertible. Then the final result follows from (16). $\square$

Given an invertible local Kan manifold X , we call $Kan(X)$ the **Kan replacement** of X .

Even though $Kan(X)$ is not Kan, its 2-truncation $\tau_2(Kan(X))$ behaves well. We define n -truncation τ_n (it is called $\tau_{\leq n}$ in [6, Section 3]), of a simplicial manifold X as,

$$\tau_n(X)_k = X_k, \forall k \leq n-1, \quad \tau_n(X)_k = X_k / \sim_k, \forall k \geq n,$$

where two elements $x \sim_k y$ in X_k if they are homotopic¹ and have the same n -skeleton. Since in the procedure, taking a quotient is involved, the result $\tau_n(X)$ might not be a simplicial manifold anymore. We view it as a simplicial stack. When X is Kan, $\tau_n(X)$ viewed as a simplicial set is always a discrete n -groupoid. It is representable, namely it is indeed a simplicial manifold, if and only if the quotient $X_n / \sim_n$ is representable because the higher levels are decided by $X_n / \sim_n$. Even though $Kan(X)$ is not a Kan manifold, we still have

Proposition 2.4. *When $Kan(X)_2 / \sim_2$ is representable, $\tau_2(Kan(X))$ is a Lie 2-groupoid.*

Proof. As a simplicial set, $Kan(X)$ is Kan. Hence $\text{hom}(\Delta[n], \tau_2(Kan(X))) \cong \text{hom}(\Lambda[n, j], \tau_2(Kan(X)))$, for $n \geq 3$. Especially, $\tau_2(Kan(X))_3 \cong \text{hom}(\Lambda[3, 0], \tau_2(Kan(X)))$. Since the higher layers are determined by the first four layers,

$$\tau_2(Kan(X)) = \text{Cosk}^3 \circ \text{Sk}^3(\tau_2(Kan(X))),$$

by the same argument in [16, Section 2.3], to show $\tau_2(Kan(X))$ is a Lie 2-groupoid, we only need to show that $\text{hom}(\Lambda[3, 0], \tau_2(Kan(X)))$ is representable and $Kan(m \leq 2, j)$ for $\tau_2(Kan(X))$. In fact the induction argument there already shows that the representability of $\text{hom}(\Lambda[3, 0], \tau_2(Kan(X)))$ is implied by $Kan(m \leq 2, j)$ for $\tau_2(Kan(X))$ given $\tau_2(Kan(X))_2 = Kan(X)_2 / \sim_2$ is representable. Hence we only need to show $Kan(m \leq 2, j)$.

As shown in Def.-Prop. 2.3, $\text{hom}(\Delta[1], X^{\beta+1}) \rightarrow \text{hom}(\Lambda[1, j], X^{\beta+1})$ being the pull-back of d_1 or d_0 , is a surjective submersion, hence $\text{hom}(\Delta[1], Kan(X)) \rightarrow \text{hom}(\Lambda[1, j], Kan(X))$ is a surjective submersion. This is $Kan(1, j)$ for $Kan(X)$, hence for $\tau_2(Kan(X))$.

The surjective part in $Kan(2, j)$ is automatically satisfied: since $Kan(X)$ is Kan as a simplicial set, the composed map

$$Kan(X)_2 \rightarrow \tau_2(Kan(X)) \xrightarrow{p} \text{hom}(\Lambda[2, j], Kan(X)) = \text{hom}(\Lambda[2, j], \tau_2(Kan(X))),$$

is surjective, hence the desired map p is also surjective. We only need to show the submersion part. Then what happened to the degenerate faces where the horn projection map is not a submersion for $Kan(X)_2$? An element $\eta \in Kan(X)_2$ can be described as a tree as stated in Lemma 3.8. If all the vertices of the tree are triangles in X_2 , then the horn projection map is a submersion for $Kan(X)_2$, hence p is a submersion. The problem happens exactly when the tree contains at least one vertex coming from one of the three copies of $X_1 \times_{X_0} X_1$. But these bad pieces as in (13) and (14) are all homotopic via elements in $Kan(X)_3$ to the boundary of good pieces as in (12), where the submersion holds. Hence the submersion part is also true for the 2-truncation. □

3. UNIVERSAL PROPERTIES

Given a local Lie 1-groupoid W (or the nerve of a local Lie groupoid), then it extends to a Lie 2-groupoid $\tau_2(Kan(W))$. In [15], we verified that $\tau_2(Kan(W))$ is always a Lie 2-groupoid (even though $\tau_1(Kan(W))$ might not be Lie) with universal property.

For this purpose, we need to show some universal properties of our Kan replacement. It should be stable under Morita equivalence of simplicial manifolds (whatever that is), and if some simplicial manifold X is already Kan, $Kan(X)$ should be Morita equivalent to X . Hence let's first begin with an introduction of these concepts such as Morita equivalence.

¹This means that $d_i x = d_i y$, $0 \leq i \leq k$, and there exists $z \in X_{k+1}$ such that $d_k(z) = x$, $d_{k+1}(z) = y$, and $d_i z = s_{k-1} d_i x = s_{k-1} d_i y$, $0 \leq i < k$.

3.1. Morita equivalence of local Kan manifolds. The reader's first guess is probably that a morphism $f : X \rightarrow Y$ of simplicial manifolds ought to be a simplicial smooth map i.e. a collection of smooth maps $f_n : X_n \rightarrow Y_n$ that commute with faces and degeneracies. We shall call such a morphism a *strict map* from X to Y . Unfortunately, it is known that, already in the case of usual Lie groupoids, such strict notions are not good enough. Indeed there are strict maps that are not invertible even though they ought to be isomorphisms. That's why people introduced the notion of *Hilsum-Skandalis bimodules* [9]. Here is an example of such a situation: consider a manifold M with an open cover $\{\mathcal{U}_\alpha\}$. The simplicial manifold X with $X_n = \bigsqcup_{\alpha_1, \dots, \alpha_n} \mathcal{U}_{\alpha_1} \cap \dots \cap \mathcal{U}_{\alpha_n}$ maps naturally to the constant simplicial manifold M . All the fibers of that map are simplices, in particular they are contractible simplicial sets. Nevertheless, that map has no inverse.

The second guess is then to define a special class of strict maps which we shall call *hypercovers*. A map from X to Y would then be a *zig-zag* of strict maps $X \xleftarrow{\sim} Z \rightarrow Y$, where the map $Z \rightarrow X$ is one of these hypercovers.

Another alternative however equivalent way to define a generalized morphism of simplicial manifolds follows from [7, Section 2.4]'s Cartesian fibrations. In this paper, we use the zig-zag method with the notion of hypercover.

Our hypercover is very much inspired from the notion of hypercover of étale simplicial objects [1, 4] and of trivial fibration of Quillen for simplicial sets [10].

Recall [8, Section I.3], given a pointed Kan simplicial set X , i.e. $X_0 = pt$, its homotopy groups are given by

$$\pi_n(X) := \{x \in X_n \mid d_i(x) = pt \text{ for all } i\} / \sim$$

where $x \sim x'$ if there exists an element $y \in X_{n+1}$ such that $d_0(y) = x$, $d_1(y) = x'$, and $d_i(y) = pt$ for all $i > 1$. When X_0 is not necessarily a point, π_n is a sheaf over X_0 in general.

Lemma 3.1. *Given a map $S \rightarrow T$ of pointed Kan simplicial sets, if for any $n \geq 0$ and any commutative solid arrow diagram*

$$(19) \quad \begin{array}{ccc} \partial\Delta[n] & \longrightarrow & S \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta[n] & \longrightarrow & T \end{array}$$

there exists a dotted arrow that makes both triangles commute, then this map is a homotopy equivalence, i.e. $\pi_n(S) = \pi_n(T)$. Here $\partial\Delta[n]$ stands for the boundary of the n -simplex.

The proof is standard.

Translating the condition of Lemma 3.1 into hom spaces gives:

Definition 3.2. A strict map $f : Z \rightarrow X$ of local Kan simplicial manifolds is a *hypercovers* if the natural map

$$(20) \quad Z_m = \text{hom}(\Delta[m], Z) \rightarrow \text{hom}(\partial\Delta[m] \rightarrow \Delta[m], Z \rightarrow X)$$

is a surjective submersion for all $0 \leq m$.

Here $\text{hom}(A \rightarrow B, Z \rightarrow X)$ denotes the pull-back spaces of the form $\text{hom}(A, Z) \times_{\text{hom}(A, X)} \text{hom}(B, X)$, where the maps are induced by some fixed maps $A \rightarrow B$ and $Z \rightarrow X$. This notation indicates that the space parameterizes all commuting diagrams of the form

$$\begin{array}{ccc} A & \longrightarrow & Z \\ \downarrow & & \downarrow \\ B & \longrightarrow & X, \end{array}$$

where we allow the horizontal arrows to vary but we fix the vertical ones.

Similarly, we can define hypercover for Lie n -groupoids:

Definition 3.3. A strict map $f : Z \rightarrow X$ of Lie n -groupoids is a *hypercover* if the natural map (20) is a surjective submersion for all $0 \leq m < n$ and is an isomorphism when $m = n$.

Remark 3.4. As proved in [14], if $f : Z \rightarrow X$ is a hypercover of Lie n -groupoids, then (20) is automatically an isomorphism for all $m > n$.

As in the case of Definition 1.1, we need to justify that the pull-back $\text{hom}(\partial\Delta[m] \rightarrow \Delta[m], Z \rightarrow X)$ is a manifold. This is rather surprising since the spaces $\text{hom}(\partial\Delta[m], Z)$ need not be manifolds (for example take $m = 2$ and Z the cross product groupoid associated to the action of S^1 on $\mathbb{R}^2$ by rotation around the origin). We justified this in [14] for Kan simplicial manifolds, but it is clear that only the submersion property is needed, hence the same proof works for local Kan manifolds.

Definition 3.5. Two local Kan simplicial manifolds X and Y are *Morita equivalent* if there is another local Kan simplicial manifold Z such that both of the maps $X \xleftarrow{\sim} Z \xrightarrow{\sim} Y$ are hypercovers. In [14, Section 2], we show that this definition does give an equivalence relation. We call it *Morita equivalence* of local Kan simplicial manifolds.

We also define *Morita equivalence* of Lie n -groupoids exactly in the same fashion using hypercover of Lie n -groupoids.

Hypercover of Lie n -groupoids may also be understood as a higher analogue of pull-back of Lie groupoids. Let X be a 2-groupoid and $Z_1 \rightrightarrows Z_0$ be two manifolds with structure maps as in (1) up to the level $n \leq 1$, and $f_n : Z_n \rightarrow X_n$ preserving the structure maps d_k^n 's and s_k^{n-1} 's for $n \leq 1$. Then $\text{hom}(\partial\Delta[n], Z)$ still makes sense for $n \leq 1$. We further suppose that $f_0 : Z_0 \rightarrow X_0$ (hence $Z_0 \times_{X_0 \times X_0} X_1$ is a manifold) and $Z_1 \rightarrow Z_0 \times_{X_0 \times X_0} X_1$ are surjective submersions. That is to say that the induced map from Z_k to the pull-back $\text{hom}(\partial\Delta[k], Z) \times_{\text{hom}(\partial\Delta[k], X)} X_k$ are surjective submersions for $k = 0, 1$. Then we form

$$Z_2 = \text{hom}(\partial\Delta[2], Z) \times_{\text{hom}(\partial\Delta[2], X)} X_2,$$

which is a manifold (see [14, Lemma 2.4]).


Moreover there are $d_i^2 : Z_2 \rightarrow Z_1$ induced by the natural projections $\text{hom}(\partial\Delta[2], Z) \rightarrow Z_1$; $s_i^1 : Z_1 \rightarrow Z_2$ by

$$s_0^1(h) = (h, h, s_0^0(d_0^1(h)), s_0^1(f_1(h))), \quad s_1^1(h) = (s_0^0(d_1^1(h)), h, h, s_1^1(f_1(h)));$$

$m_i : \text{hom}(\Lambda[3, i], Z) \rightarrow Z_2$ by for example

$$m_0((h_2, h_5, h_3, \bar{\eta}_1), (h_4, h_5, h_0, \bar{\eta}_2), (h_1, h_3, h_0, \bar{\eta}_3)) = (h_2, h_4, h_1, m_0(\bar{\eta}_1, \bar{\eta}_2, \bar{\eta}_3)),$$

and similarly for other m 's.



Then $Z_2 \rightrightarrows Z_1 \rightrightarrows Z_0$ is a Lie 2-groupoid and we call it the *pull-back 2-groupoid* by f . Moreover $f : Z \rightarrow X$ is an equivalence with the natural projection $f_2 : Z_2 \rightarrow X_2$.

3.2. Lemmas. What we wish to prove is: If X is already a Kan simplicial manifold, then $X \xrightarrow{\sim} Kan(X)$ are Morita equivalent. It is very easy to prove for simplicial set. Since the procedure of Kan replacement is basically to fill out horns, the geometric realization of $Kan(X)$ and X are homotopic to each other. Since X is Kan, this is equivalent to (19). However, the missing tool of homotopy theory of simplicial manifolds (which do not form a model category, but building a certain machinery as a suitable replacement of model category should be the eventual correct method to prove these lemmas.) prevents us to apply this proof directly. In fact, in the case of simplicial sets, one can easily obtain a morphism $\pi : Kan(X) \rightarrow X$ such that the composition $X \rightarrow Kan(X) \rightarrow X$ is the identity. Then it is straightforward to check that $Kan(X) \rightarrow X$ has the correct lifting property. However, π is not unique (basically it depends on the choice of fillings in the Kan condition). Hence when generalized to a differential category, π is in general not a continuous morphism. This forces us to use another proof. Here we provide a proof for Lie 2-groupoids W .

Lemma 3.6. *If W is a Lie 2-groupoid, then $\tau_2(Kan(W))$ is a Lie 2-groupoid which is Morita equivalent to W .*

Proof. Usually, we do not have a direct map from $\tau_2(Kan(W))$ to W because there is no (unique) multiplication map $W_1 \times_{W_0} W_1 \rightarrow W_1$ (even when there exists such a multiplication, we will encounter the issue of surjective submersions). Hence we must construct a middle step.


A more natural way to describe this is to use the corresponding stacky groupoid $\mathcal{G} \Rightarrow W_0$, where $\mathcal{G}$ is presented by the Lie groupoid $G_1 \Rightarrow G_0$, with $G_0 = W_1$ and G_1 the set of bigons in W_2 , and the multiplication $\mathcal{G} \times_{W_0} \mathcal{G} \rightarrow \mathcal{G}$ is presented by bimodule $E_m = W_2$. The bimodules of various compositions of multiplication from various copies of $\mathcal{G}$ to $\mathcal{G}$ are presented by various fibre product of W_2 's. For example, the bimodule $W_2 \times_{d_1, W_1, d_2} W_2$ with the moment map J_l to $W_1 \times_{W_0} W_1 \times_{W_0} W_1$ and J_r to W_1 , presents the multiplication

$$m \circ (m \times id) : (\mathcal{G} \times_{W_0} \mathcal{G}) \times_{W_0} \mathcal{G} \rightarrow \mathcal{G}.$$

To simplify the notation, we denote a k -times fibre product as $\square^{\times k}$ when it's clear from the context. We construct $Z_0 = W_0 = W_0$ and, Z_1 is the disjoint union of these bibundles $W_2^{\times k}$ presenting different compositions of multiplication,

$$Z_1 = W_1 \sqcup W_2 \sqcup (W_2 \sqcup W_2^{\times 2} \sqcup W_2^{\times 2} \sqcup W_2^{\times 3}) \sqcup \dots$$

It is best to be understood as the following picture:



That is, we fill out horns in $Kan(W)_1$ by replacing $W_1^{\times n}$ with $W_2^{\times(n-1)}$. The projections of $Z_1 \rightarrow Kan(W)_1$ and $Z_1 \rightarrow W_1$ are simply the disjoint union of the left and right moment maps respectively. These projections are both surjective submersions.

To show that $\tau_2(Kan(W))$ is Morita equivalent to W , we only have to show that the pullback 2-groupoids on Z are the same, that is

$$(21) \quad Kan(W)_2 / \sim_2 \times_{\text{hom}(\partial\Delta[2], Kan(W))} \text{hom}(\partial\Delta[2], Z) \cong W_2 \times_{\text{hom}(\partial\Delta[2], W)} \text{hom}(\partial\Delta[2], Z).$$

If the map $p : M \rightarrow N$ is surjective and admit local section at any point in N , then the pull-back groupoid $G_1 \times_M N \Rightarrow G_0 \times_M N$ is free and proper if and only the original groupoid $G_1 \Rightarrow G_0$ is so. Since this is our case, the isomorphism (21) automatically implies that $Kan(W)/\sim_2$ is representable. By Prop. 2.4, $\tau_2(Kan(W))$ is a Lie 2-groupoid.

We denote the two pullbacks by the map $Z_1 \rightarrow W_1$ and $Z_1 \rightarrow Kan(W)_1$ to Z_1 by $W|_Z$ and $Kan(W)|_Z$ respectively, and we construct morphisms

$$\pi : (Kan(W)|_Z)_2 \rightarrow (W|_Z)_2, \quad \iota : (W|_Z)_2 \rightarrow (Kan(W)|_Z)_2,$$

and prove $\pi \circ \iota = id$ and $\iota \circ \pi \sim id$ up to something in $(Kan(W)|_Z)_3$. Then the above isomorphism follows naturally. Notice that $Kan(W)$ is not a Lie 2-groupoid usually, but pull-back described in Section 3.1 works also when X is a local Kan manifold. We form $(X|_Z)_n = \text{hom}(sk_1(\Delta[n]) \rightarrow \Delta[n], Z \rightarrow X)$, where sk_1 denotes of taking the 1-dimensional skeleton. By [14, Lemma 2.4], $(X|_Z)_n$ are manifolds. Then it's easy to check that $\tau_2(Kan(W)|_Z) = \tau_2(Kan(W))|_Z$.

We first construct ι . Let S be a simplicial polygon with three marked points, namely a simplicial set constructed inductively

$$\Delta[2] = S_0 \hookrightarrow S_1 \hookrightarrow S_2 \hookrightarrow \dots \hookrightarrow S_i \dots,$$

by push-out $S_{i+1} = S_i \sqcup_{\Delta[1]} \Delta[2]$ and the three marked points are the vertices of S_0 . With these three marked points, the S_i 's can be viewed as generalized triangles with their three sides a concatenation of line segments. In this sense, we also have the three facial maps d^k .

We have a natural embedding $W \hookrightarrow Kan(W)$, but this embedding does not give $W|_Z \rightarrow Kan(W)|_Z$. In fact, take an element $(w, \partial z) \in (W|_Z)_2 = \sqcup_i \text{hom}(S_i, W)$ for a certain set of S_i 's, then $(w, \partial z) \notin (Kan(W)|_Z)_2$ since ∂w the boundary of w , is not ∂z under the map $\partial_2 Z \rightarrow \partial_2 Kan(W)$. Here $\partial_k \square = \text{hom}(\partial \Delta[k], \square)$. To construct ι we need to construct a morphism $\mu_i : \text{hom}(S_i, W) \rightarrow Kan(W)_2$ inductively, such that it commutes with the facial map d_k for $k = 0, 1, 2$,

$$(22) \quad \begin{array}{ccc} \text{hom}(S_i, W) & \xrightarrow{d_k} & \text{hom}(d^k S_i, W) \\ \downarrow \mu_i & \nearrow \partial & \\ Kan(W)_2 & & \end{array}$$


Then $\iota(w, \partial z) := (\mu_i(w), \partial z)$ where $w \in \text{hom}(S_i, W)$.

Step 1: We first prove the case $i = 1$. We simplify the notation by $K := Kan(W)$.

$$\begin{aligned}
 (23) \quad & \text{hom}(S_1, W) = W_2 \times_{d_k, W_1, d_1} W_2 \\
 & \downarrow \text{by } W_1 \times_{W_0} W_1 \hookrightarrow K_2 \\
 & W_2 \times_{d_k, W_1, d_1} W_2 \times_{W_1 \times_{W_0} W_1} K_2 \\
 & \downarrow \text{by } W \hookrightarrow K \\
 & \text{hom}(\Lambda[3, k'], K) \\
 & \downarrow Kan!(3, k') \\
 & \text{hom}(\Delta[3], K) \\
 & \downarrow d_{k'} \\
 & K_2
 \end{aligned}$$


But this map does not commute with the facial map (see (22)). For this purpose, we only need to compose with the following one,

$$\begin{aligned}
 (24) \quad & K_2 \rightarrow (K_2 \times_{W_1} (W_1 \times_{W_0} W_1)) \times_{K_1 \times_{W_0} W_1} W_1 \times_{W_0} W_1 \\
 & \downarrow W_1 \times_{W_0} W_1 \hookrightarrow K_2 \\
 & \text{hom}(\Lambda[3, k''], K) \rightarrow K_2
 \end{aligned}$$



Step 2: Now suppose we have such a map $\text{hom}(S_i, W) \rightarrow K$, then we can construct a map $\text{hom}(S_{i+1}, W) \rightarrow K$ as below,

$$\begin{array}{ccc}
 \text{hom}(S_{i+1}, W) = \text{hom}(S_i, W) \times_{W_1} W_2 & \xrightarrow{\partial} & \text{hom}(\partial S_{i+1}, W) = \text{hom}(\partial S_i, W) \times_{W_1} \partial_2 W \\
 \downarrow \text{By Lemma 3.7} & & \downarrow \\
 \text{hom}(S_i, W) \times_{d_k, K_1, d_1} K_2 & \xrightarrow{\partial} & \text{hom}(\partial S_i, W) \times_{K_1} \partial_2 K \\
 \downarrow \text{By } \text{hom}(S_i, W) \rightarrow K_2 & & \downarrow \\
 K_2 \times_{d_k, K_1, d_1} K_2 = \text{hom}(S_1, K) & \xrightarrow{\partial} & \text{hom}(\partial S_1, K) \\
 \downarrow \text{Similarly as Step 1, replace } W \text{ by } K & \nearrow \partial & \\
 K_2 & &
 \end{array}$$



Lemma 3.7. *There is a natural morphism $K_1 \times_{W_1} W_2 \rightarrow K_2$.*

We leave this to the readers as an exercise involving the Kan condition of K .

To construct π , we first construct a local morphism $f : Kan(W) \rightarrow W$ inductively. The first step is to construct f_n^1 by

$$\begin{aligned} W_n^1 &= \text{hom} \left(\Delta[n], W \coprod_{\coprod \Lambda[k,j] \times \text{hom}(\Lambda[k,j], W)} \coprod \Delta[k] \times \text{hom}(\Lambda[k,j], W) \right) \\ &\rightarrow W_n \coprod \text{hom}(\Delta[n], \Delta[k]) \times \text{hom}(\Delta[k], W) \\ &\quad \dots \\ &\rightarrow W_n. \end{aligned}$$


In the second last step we use the strict Kan condition $\text{hom}(\Lambda[k,j], W) \cong W_k$ when $k \geq 2$ and we choose a local section $\text{hom}(\Lambda[2,1], W) \rightarrow W_2$ when $k = 2$. The last step follows from the composition $\text{hom}(\Delta[n], \Delta[k]) \times \text{hom}(\Delta[k], W) \rightarrow W_n$ and thus both spaces in the coproduct have a natural map to W_n .

Suppose that $f^\beta : W^\beta \rightarrow W$ is constructed. Then $f^{\beta+1}$ is the composition of the following natural morphisms

$$\begin{aligned} W_n^{\beta+1} &= \text{hom} \left(\Delta[n], W^\beta \coprod_{\coprod \Lambda[k,j] \times \text{hom}(\Lambda[k,j], W^\beta)} \coprod \Delta[k] \times \text{hom}(\Lambda[k,j], W^\beta) \right) \\ &\rightarrow \text{hom} \left(\Delta[n], W \coprod_{\coprod \Lambda[k,j] \times \text{hom}(\Lambda[k,j], W)} \coprod \Delta[k] \times \text{hom}(\Lambda[k,j], W) \right) \\ &= W_n^1 \xrightarrow{f_n^1} W_n. \end{aligned}$$

Then f is the colimit of f^β .

More geometrically, if we view an element in $Kan(W)_2$ as a set of small triangles of W_2 touching together, f_2 is basically to compose these small triangles into a big one in W_2 with a choice of filling for each $W_1 \times_{W_0} W_1$, which is given by f^1 .



Now when we make a choice of fillings for a $W_1 \times_{W_0} W_1$ on the boundary, instead of choosing some filling given by f^1 , we choose the element $\partial z \in \text{hom}(\partial\Delta[2], Z)$, then this element in W_2 is denoted by $f_2(x) \circ \partial z$.

Thus f_2 induces a map

$$(25) \quad Kan(W)_2 \times_{\text{hom}(\partial\Delta[2], Kan(W))} \text{hom}(\partial\Delta[2], Z) \xrightarrow{\pi} W_2 \times_{\text{hom}(\partial\Delta[2], W)} \text{hom}(\partial\Delta[2], Z).$$


as $(x, \partial z) \mapsto (f_2(x) \circ \partial z, \partial z)$. In Lemma 3.8, we give a combinatorial proof that this map does not depend on the choice of fillings. Hence we obtain a well-defined global map π .

Then it is not hard to see that $\pi \circ \iota = id$ since π is exactly the opposite procedure of ι .

The procedure to form π and ι is basically to use $Kan!(3, j)$ to compose (for example (23)), hence $\iota \circ \pi$ and id differ by something in $Kan(W)_3$. $\square$

Lemma 3.8. *The map π does not depend on the choice of sections in the construction of f .*

Proof. We denote an element in $\eta \in Kan(W)_2$ by a bicolored tree




A point is black if it represents a weird triangle, i.e. a triangle comes from the first copy of $X_1 \times_{X_0} X_1$ which serves as an artificial filling; otherwise, it is white. For the other two copies of $X_1 \times_{X_0} X_1$, they are degenerate ones and can not glue directly with a normal triangle in W_2 . Since degenerate elements play the role of identities in composition π , we here ignore them. We prove the result by induction on the number of generations and the number of points in the youngest generation. It is obvious for the initial case.

Now take three siblings points in the youngest generation, if all of them are white, then we use $Kan(3, j)$ without a choice and we end up with an element $\eta' \in Kan(W)_2$ which has a fewer number of generations or a fewer number of points in the youngest generation. Done!

If one of the three siblings is black, then there is precisely one black one in these three siblings, which we denote by x . Since $\text{hom}(\partial\Delta[2], Z)$ will give the fillings for the weird triangles on the border of η , to show the independence, we only have to deal with the inner triangles. Then some ancestor of x must have a black descendant y , because a weird triangle must lie on the side of another triangle (which is the parent of y).

The simplest situation in this case is when the other black descendant is a (true) cousin (namely their direct ancestors are siblings)


(26)



We choose a triangle η_{034} , and we are given triangle η_{014} , η_{134} , η_{123} , η_{234} , η_{024} . The procedure is to compose η_{034} , η_{014} , η_{134} first to obtain η_{013} by $Kan(3, 3)$; then to compose

η_{034} , η_{024} , η_{234} secondly to obtain η_{023} by $Kan(3,3)$; finally to compose η_{013} , η_{023} , and η_{123} to obtain η_{012} by $Kan(3,3)$. These can be viewed as multiplications for 2-groupoid [14, Section 2.3]. By associativity of such multiplications (or equivalent $Kan(3,j)!$ and $Kan(4,j)!$), we can obtain the same η_{012} by another order of composition, namely we use $Kan(3,2)$ first to obtain η_{124} then $Kan(3,3)$. Since the second way to compose does not depend on the choice of η_{034} , our final result η_{012} does not depend on the choice either. Hence by the induction hypothesis, we will choice-independently end up with an element $f_2(x) \circ \partial z$.

We might meet more complicated situations, namely the other black descendant y is a more remote cousin, but we can reduce them to the simple situation above:



We choose a triangle $\xi \in W_2$ corresponding to x in the youngest generation. Then we use $Kan(3,j)$ to compose to obtain ξ' and ξ'' corresponding to x' and y' respectively. As shown in the picture, the true cousins x', y' are ancestors of x and y respectively. We also do the same for the other branch and obtain ζ . During this procedure, we might have to make other choices of fillings for other black points. But it does not matter, since our point is to show the independence on the choice ξ . Then we are again in a similar situation as of (26). We choose a filling η_{234} . By $Kan(4,j)!$ the final result does not depend on the order of composition. Then by $Kan(3,2)$ we first obtain η_{124} ; by the induction hypothesis, we obtain η_{024} independent of the choice of ξ because η_{024} contains fewer descendants; finally, by $Kan(3,3)$ we obtain η_{012} , which is independent of the choice of filling ξ . $\square$

Lemma 3.9. *If both X and Y are Lie n -groupoids, then a hypercover of local Kan simplicial manifolds $X \xrightarrow{\sim} Y$ is automatically a hypercover of Lie n -groupoids.*

Proof. Since X is a Lie n -groupoid, we have the composed morphisms $\text{hom}(\Lambda[n+1, j], X) \cong X_{n+1} \xrightarrow{d_j} X_n \rightarrow Y_n$ and $\text{hom}(\Lambda[n+1, j], X) \cong X_{n+1} \xrightarrow{d_j} X_n \xrightarrow{\partial} \text{hom}(\partial\Delta[n], X)$. This gives us a map $\text{hom}(\Lambda[n+1, j], X) \xrightarrow{p} \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)$. With this map, we rewrite

$$\text{hom}(\partial\Delta[n+1] \rightarrow \Delta[n+1], X \rightarrow Y) \cong \text{hom}(\Lambda[n+1, j], X) \times_{\text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)} X_n.$$

Since $X \xrightarrow{\sim} Y$ as local Kan simplicial manifolds, the following map

$$X_{n+1} \cong \text{hom}(\Lambda[n+1, j], X) \rightarrow \text{hom}(\Lambda[n+1, j], X) \times_{\text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)} X_n$$

is a surjective submersion. This implies that $X_n \rightarrow \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)$ is injective. However, $X_n \rightarrow \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)$ is a surjective submersion by the condition of hypercovers. Hence $X_n \cong \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], X \rightarrow Y)$, which shows $X \xrightarrow{\sim} Y$ as Lie n -groupoids. $\square$

This implies

Corollary 3.10. *Two Lie n -groupoids $X \xrightarrow{\sim} Y$ are Morita equivalent as local Kan simplicial manifolds if and only if they are Morita equivalent as Lie n -groupoids.*

Lemma 3.11. *If $\phi : X \xrightarrow{\sim} Y$ is a hypercover of local Kan simplicial manifolds, and if $\text{Kan}(X)_2 / \sim_2$ is representable, then both $\tau_2(\text{Kan}(X))$ and $\tau_2(\text{Kan}(Y))$ are Lie 2-groupoids and the induced map $\tau_2(\text{Kan}(X)) \rightarrow \tau_2(\text{Kan}(Y))$ is a hypercover of Lie 2-groupoids.*

Proof. We first show that if $\phi : K \rightarrow K'$ is a hypercover of Kan simplicial sets (i.e. (20) is surjective instead of a surjective submersion), then the natural map

$$(27) \quad \tau_n(K)_n \xrightarrow{f} \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], \tau_n(K) \rightarrow \tau_n(K')),$$

is an isomorphism. Notice that the right hand side is simply $\text{hom}(\partial\Delta[n], K) \times_{\text{hom}(\partial\Delta[n], K')} \text{hom}(\Delta[n], \tau_n(K'))$. Thus we have a commutative diagram

$$\begin{array}{ccc} K_n & \xrightarrow{\quad \twoheadrightarrow \quad} & \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], K \rightarrow K') \\ g \downarrow & & \downarrow \\ \tau_n(K)_n & \xrightarrow{\quad f \quad} & \text{hom}(\partial\Delta[n] \rightarrow \Delta[n], \tau_n(K) \rightarrow \tau_n(K')), \end{array}$$

where $\twoheadrightarrow$ denotes surjective maps. Then f must be surjective because $f \circ g$ being surjective implies f being surjective.

Take $(\delta x_n, [y_n]) \in \text{hom}(\partial\Delta[n], K) \times_{\text{hom}(\partial\Delta[n], K')} \text{hom}(\Delta[n], \tau_n(K'))$. If both $[x_n], [x'_n] \in \tau_n(K)_n$ map to $(\delta x_n, [y_n])$, that is the boundary $\partial x_n = \partial x'_n = \delta x_n \in \text{hom}(\partial\Delta[n], K)$ and $\phi_n(x'_n) = y'_n \sim y_n \sim y''_n = \phi_n(x_n)$, then y''_n and y'_n differ by a certain element $y_{n+1} \in K'_{n+1}$. Since

$$K_{n+1} \twoheadrightarrow \text{hom}(\partial\Delta[n+1] \rightarrow \Delta[n+1], K \rightarrow K'),$$

is surjective, there exists x_{n+1} such that $\phi_{n+1}(x_{n+1}) = y_{n+1}$ and x_n, x'_n differ by x_{n+1} . This proves that $[x_n] = [x'_n] \in \tau_n(K)_n$. Hence f is also injective.

If the map $p : M \rightarrow N$ is surjective and admits local section at any point in N , then the pull-back groupoid $G_1 \times_M N \rightrightarrows G_0 \times_M N$ is free and proper if and only the original groupoid $G_1 \rightrightarrows G_0$ is so. Since this is our case, the isomorphism (27), when applied to $n = 2$ and $K = \text{Kan}(X)$ $K' = \text{Kan}(Y)$, implies that $\text{Kan}(Y) / \sim_2$ is representable. Hence $\tau_2(\text{Kan}(X))$ and $\tau_2(\text{Kan}(Y))$ are Lie 2-groupoids by Prop. 2.4.

Now we only need to verify that the morphism

$$\tau_2(\text{Kan}(X))_m \rightarrow \text{hom}(\partial\Delta[m] \rightarrow \Delta[m], \tau_2(\text{Kan}(X)) \rightarrow \tau_2(\text{Kan}(Y)))$$

is a surjective submersion for $m = 0, 1$. For $m = 0$ it is implied by $X_0 \rightarrow Y_0$ being a surjective submersion. For $m = 1$, by induction, we need to show that the natural map

$$(28) \quad X_1^{\beta+1} \rightarrow \text{hom}(\partial\Delta[1] \rightarrow \Delta[1], X^{\beta+1} \rightarrow Y^{\beta+1}),$$

is a surjective submersion supposing the same is true for β . We have

$$X_1^{\beta+1} = X_1^\beta \sqcup \text{hom}(\Lambda[2, 1], X^\beta), \quad Y_1^{\beta+1} = Y_1^\beta \sqcup \text{hom}(\Lambda[2, 1], Y^\beta).$$

The right hand side of (28) decomposes into two terms I, II according to the decomposition of $Y_1^{\beta+1}$,

$$\begin{aligned} I &= \text{hom}(\partial\Delta[1], X^\beta) \times_{\text{hom}(\partial\Delta[1], Y^\beta)} \text{hom}(\Delta[1], Y^\beta) \\ II &= \text{hom}(\partial\Delta[1], X^\beta) \times_{\text{hom}(\partial\Delta[1], Y^\beta)} \text{hom}(\Lambda[2, 1], Y^\beta) \end{aligned}$$



By the induction hypothesis, $X_1^\beta \rightarrow I$ is a surjective submersion. Further by [14, Lemma 2.5] (take $S = T = \Lambda[2, 1]$, and $T' = \partial\Delta[1]$),




$$\mathrm{hom}(\Lambda[2, 1], X^\beta) \rightarrow II$$



is a surjective submersion. Thus (28) is a surjective submersion. $\square$


REFERENCES

- [1] *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*. Lecture Notes in Mathematics, Vol. 269. Springer-Verlag, Berlin, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat.
- [2] A. S. Cattaneo and G. Felder. Poisson sigma models and symplectic groupoids. In *Quantization of singular symplectic quotients*, volume 198 of *Progr. Math.*, pages 61–93. Birkhäuser, Basel, 2001.
- [3] M. Crainic and R. L. Fernandes. Integrability of Lie brackets. *Ann. of Math. (2)*, 157(2):575–620, 2003.
- [4] E. M. Friedlander. *Étale homotopy of simplicial schemes*, volume 104 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, N.J., 1982.
- [5] E. Getzler. Lie theory for nilpotent L-infinity algebras, arxiv:math.AT/0404003.
- [6] A. Henriques. Integrating L_∞ -algebras. *Compos. Math.*, 144(4):1017–1045, 2008.
- [7] J. Lurie. Higher Topos Theory, arXiv:math/0608040v4 [math.CT].
- [8] J. P. May. *Simplicial objects in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992. Reprint of the 1967 original.
- [9] J. Mrčun. *Stability and invariants of Hilsum-Skandalis maps*. Dissertation, Utrecht University, Utrecht, 1996.
- [10] D. G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [11] H.-H. Tseng and C. Zhu. Integrating Lie algebroids via stacks. *Compos. Math.*, 142(1):251–270, 2006.
- [12] W. T. van Est and M. A. M. van der Lee. Enlargeability of local groups according to Mal’cev and Cartan-Smith. In *Action hamiltoniennes de groupes. Troisième théorème de Lie (Lyon, 1986)*, volume 27 of *Travaux en Cours*, pages 97–127. Hermann, Paris, 1988.
- [13] P. Ševera. L infinity algebras as 1-jets of simplicial manifolds (and a bit beyond) , arXiv:math/0612349 [math.DG].
- [14] C. Zhu. n-groupoids and stacky groupoids, arxiv:math.DG/0801.2057[math.DG].
- [15] C. Zhu. Lie II theorem for Lie algebroids via stacky Lie groupoids, arxiv:math/0701024 [math.DG].
- [16] C. Zhu. Lie n-groupoids and stacky Lie groupoids, arxiv:math.DG/0609420.









X2

$\Lambda^{1,0}$

$\Lambda^{1,1}$

$\Lambda^{2,0}$

$\Lambda^{2,1}$

$\Lambda^{2,2}$

$\Lambda^{3,0}$

$\Lambda^{3,1}$

$\Lambda^{3,2}$

$\Lambda^{3,3}$




$\Lambda^{4,0}$

$\Lambda^{4,1}$

$\Lambda^{4,2}$

$\Lambda^{4,3}$

$\Lambda^{4,4}$



...

